



Ecological Organic Agriculture initiative for Africa SDC- BvAT (EOA-Phase II) project Pillar I. Research Result

ON

Performance of carrot(*Daucus carota*) under different fertilizer types in Tehuledere District, South Wollo Zone , Eastern Amhara, Ethiopia

BY

Name	Academic rank	Email
Dr. Faris Hailu	Associate professor	markhmets@yahoo.com
Mr. Seid Hassen	Assistant professor	seidhassen19@yahoo.com
Dr. Eshetu Belete	Assistant professor	ebelete2022@yahoo.com
Dr. Seid Hussen	Assistant professor	hamidashm@gmail.com
Tewodros Alemu	Assistant professor	teda2010@gmail.com

Wollo University, Dessie, POBox-1145, Ethiopia

ABSTRACT

The demand for organic agricultural practices has been growing due to concerns about the environmental and health impacts of chemical-intensive farming. This study aimed to investigate the effects of various organic fertilizers on carrot (Nantes) growth and yield in the specific agro-climatic conditions of Tehuledere district. The experiment was conducted at two sites (Gobeya and Libannos) and using five fertilizer types (vermicompost, conventional compost, mixture of the two at 1:1 ratio, inorganic fertilizer (NPSB), and control). The experiment was laid out by using randomized complete block design with three replications, and the collected data were analysed using R-software. The study analyzed various phenological and growth parameters (emergence percentage, plant height, leaf number, fresh weight, and dry weight of leaves), and yield and yield-related parameters (root length, root diameter, fresh weight of roots, marketable root yield, unmarketable root yield, and total root yield). The results demonstrated that application of fertilizers in general significantly improved ($p < 0.001$) growth and yield compared to the control. Applications of vermicompost solely and NPSB showed the highest values ($p < 0.001$) for most of the tested parameters. Marketable and total yields were $41.7\text{t/ha} \pm 0.76$ and $45.5\text{t/ha} \pm 0.9$, and $42.5\text{t/ha} \pm 1.14$ and $45.6\text{t/ha} \pm 0.9$ for sole vermicompost and NPSB, respectively. Sole vermicompost and NPSB produced similar results ($p > 0.001$); however, NPSB poses negative environmental impact that could not result in sustainability. Therefore, organic carrot production using sole application of vermicompost is recommended for farmers of Tehuledere district and similar agroecological regions to enhance productivity while contributing to sustainable agriculture. These results have implications for policymakers, researchers, and farmers interested in promoting environmentally friendly and sustainable agricultural practices in Ethiopia.

Keywords: Organic fertilizer, Inorganic fertilizer, Carrot, Sustainable agriculture, Tehuledere

INTRODUCTION

Carrot (*Daucus carota* L.) is one of the most widely consumed root vegetables worldwide, renowned for its nutritional value and versatility. This vegetable contains phytochemicals, vitamins and pro-vitamins that prevent chronic diseases, such as cancer, cardiovascular disease, and diabetes (Nambia et al., 2010; Jamuna et al., 2011). In recent years, there has been a growing interest in organic agricultural practices, driven by concerns about the environmental and health impacts of conventional chemical-intensive farming. As a result, research into ecological organic carrot production has gained significant attention, aiming to develop sustainable and environmentally friendly approaches to enhance crop productivity. This proposed study seeks to investigate the effects of various organic fertilizer types on carrot growth, yield, and quality in the specific agro-climatic conditions of Tehuledere District. By examining the performance of carrot crops under different organic fertilizer treatments, this research aims to contribute valuable insights to sustainable agricultural practices, specifically in organic carrot production.

Ethiopia, with its diverse agro-climatic conditions, provides a favorable environment for carrot cultivation. However, traditional agricultural practices in the country have heavily relied on synthetic fertilizers, pesticides, and herbicides, leading to soil degradation, water pollution, and negative impacts on human health. Consequently, there is a pressing need to transition towards more sustainable and eco-friendly farming methods. In recent years, several studies have explored the potential of organic farming as an alternative approach to improve agricultural sustainability. Organic fertilizers, such as compost, manure, and green manure, are considered essential components of organic agriculture, as they enhance soil fertility, increase nutrient availability, and promote beneficial soil microorganisms. Studies by Basel and Sami (2014) and Shashi et al. (2018) have demonstrated the positive effects of organic fertilizers on crop growth and yield in different regions of Ethiopia.

However, specific research on organic carrot production in Tehuledere District remains limited, and there is a knowledge gap regarding the comparative performance of different organic fertilizers on carrot crops in this region. Understanding the effectiveness of various organic fertilizer types on carrot growth in the specific agro-climatic conditions of Tehuledere District is crucial for farmers to adopt sustainable agricultural practices and improve their livelihoods.

The rationale behind this study is to address the challenges posed by conventional agricultural practices in Tehuledere District and explore sustainable alternatives for carrot production. By evaluating the performance of carrot crops under different organic fertilizer types, this research aims to assess the growth and yield response of carrot crops to various fertilizers commonly available in the area, and to investigate the economic feasibility and practicality of adopting organic carrot production methods for local farmers in Tehuledere District.

The primary research problem addressed in this study is the lack of comprehensive data on the performance of carrot cultivation using organic fertilizers in Tehuledere District. Consequently, farmers may be hesitant to shift from conventional to organic practices due to uncertainty about its efficacy and economic viability.

The main objective of this research is to evaluate the performance of carrot crops under different types of fertilizers in Tehuledere District, Ethiopia. By addressing this objective, the study endeavors to contribute to the body of knowledge on ecological organic carrot production and offer valuable insights to farmers, policymakers, and researchers interested in sustainable agriculture in Ethiopia. To achieve the objective, the study aims to evaluate the growth and yield of carrot under various fertilizer types, and to identify the best fertilizer type for optimum yield and sustainable organic carrot production in Tehuledere district and areas with similar agroecology.

MATERIAL AND METHODS

Study Area

The experiment was conducted in Tehuledere district at two sites (Gobeya and Libannos) in 2022/2023 cropping season. The district is characterized by ago-ecologically moist midland.

Seed Material

The kind of carrot seed used in the experiment was Nantes and was bought from a seed importer company residing in Addis Ababa. Before planting, the seeds were first checked for their germination rate. According to Getachew and Mohammed (2012), Nantes has orange color and cylindrical roots with a blunt end and strong leaves. Farmers are highly needed for its good

adaptation in highlands (1600-2400m and annual precipitation 760-1010mm), for its high market demand, and for its good color, thick and long roots and sweet taste.

Fertilizer Type and Experimental Design

This experiment evaluated different organic fertilizers for carrot production at Gobeya and Libannos (Haiké Estifanos) in Tehuledere district. Five treatments were used (T₁: Vermicompost, T₂: Conventional compost, T₃: Combination of vermicompost 50% and 50% conventional compost, T₄: inorganic fertilizer /NPSB/, and T₅: control) and were replicated three times using Randomized Complete Block Design. Vermicompost and conventional compost rates were 3 t/ha and 12 t/ha, respectively. NPSB Fertilizer was used based on recommended rate. A plot size of 3 m by 3 m (9 m²) with a spacing of 1 m between blocks and 0.5 m between plots was used.

Experimental Field Management

The land was ploughed three times using oxen. All treatments were applied on well prepared experimental plots. The carrot seeds were drilled on experimental plots with recommended row spacing. After sowing, all the necessary agronomic management practices were done whenever required to keep the plants in healthy growth.

Data Type and Data Collection Technique

Phonological and vegetative growth parameters

Emergence percentage (%): After emergence of first seedling of every treatment, the numbers of emerged seedlings were counted daily up to 14 days after sowing. Emergence percentage was

calculated as: Emergence (%) = $\frac{\text{total number of emerged seedling}}{\text{total seeds sown}} * 100$

Plant height (cm): Plant heights from the ground level to the tip of upper most part of 10 randomly taken plants in net plot area was measured at maturity using meter tape and the mean values was calculated and used for further analysis.

Leaf number: The number of leaves from ten randomly taken plants in the net plot area was counted at the time when plants fully covered the ground surface and mean values was calculated and used for further analysis.

Fresh weight of leaves per plant (mg): It was recorded by weighing the fresh weight of ten randomly selected plants from net plot area using sensitive balance, and then the average value was computed and used for further analysis.

Dry weight of leaves per plant (mg): Leaves from ten randomly taken carrot leaves was cut in to small pieces and oven dried. After oven drying, the samples were weighed by an electrical balance and the dry matter content was calculated and used for further analysis.

Yield and yield related parameters

Root length (cm): The average length of ten randomly taken roots was recorded by a meter scale from the point of attachment of the leaves (proximal end) to the last point of the root (distal end) in each treatment combination.

Diameter of root (cm): It was recorded by measuring the root diameter about two centimeter below the root collar of ten randomly selected carrot roots harvested from the net plot area using caliper and the mean value was computed and used for further analysis.

Fresh weight of root per plant (g/root): It was recorded by weighing ten randomly selected carrot roots harvested from the net plot using sensitive balance and the mean value was computed and used for further analysis.

Root dry weight (g): Immediately after harvest, roots were cleaned thoroughly by washing with water and air dried. Then ten randomly taken carrot roots was cut into small pieces and sun dried for 3 days and then oven dried for 72 hours at 70-80 °C temperature. After oven drying, the samples were weighted by an electrical balance and dry matter content was calculated.

Marketable root yield ($t\ ha^{-1}$): Carrot roots, which are free from mechanical damages, disease and insect pest attack, small sizes (<50 g) and cracks, was considered as marketable. The weight of such carrots harvested from the net plot area was weighed using scale and expressed as ton per hectare.

Unmarketable root yield ($t\ ha^{-1}$): Carrot roots which are diseased, insect pest damaged, cracked and under sized (<50g) was considered as unmarketable. The weight of such carrots harvested from the net plot area was weighed using scale and expressed as t/ha.

Total root yield ($t\ ha^{-1}$): The total root yield was calculated by converting the yield obtained from plots to hectare and expressed as t/ha.

Statistical Analysis

Data was subjected to analysis of variance using R-software. The treatment means were compared using Tukey test at 5% level of significance.

RESULTS AND DISCUSSION

The results and discussion of growth, and yield and yield related parameters of carrot (Nantes) produced over different fertilizer types and locations are indicated below.

Growth Parameters of Carrot under Different Fertilizers and Locations

Table 1 and 2 presents the results of a study comparing the effects of different fertilizer types (vermi-compost, conventional compost, compost mixture (conventional compost and vermi-compost at 1:1 ratio), and NPSB inorganic fertilizer) and the control on the vegetative growth parameters (plant height, emergence percentage, leaf number, fresh weight of leaves per plant, and dry weight of leaves per plant) of carrot under two different environments. The data of each parameter is presented as means \pm standard error and range.

The use of organic fertilizers (vermi-compost, conventional compost, and the mixture) had a significant effect ($p < 0.001$) on the plant height, emergence percentage, leaf number, fresh weight of carrot leaves, and dry weight of carrot leaves compared to the control. However, the mean values of the parameters of the organic fertilizers were similar ($p > 0.05$) with the results of the inorganic fertilizer (NPSB). The use of NPSB, while providing comparable result, may result in negative impacts on soil health and environmental sustainability (Alzamel et al., 2022). So, organic fertilizers could be viable alternatives to inorganic fertilizers for improving carrot growth and yield. This finding was in line with other studies that have reported the positive effects of organic fertilizers on plant growth and health (Hati and Bandyopadhyay, 2011; Tadila, and Amare, 2019). The study also indicated that there was no significant interaction ($p > 0.05$) between treatments and the localities for the emergence percentage, leaf number, and fresh weight of carrot leaves. However, there was a significant interaction for the plant height ($p < 0.01$) and dry weight ($p < 0.05$) of carrot leaves (Table 1 and 2).

Height and emergence rate of carrot

Table 1 presents the results of the experiment, evaluating the plant height and emergence percentage of carrot under different fertilizer treatments in Tehuledere district. The data is presented for five fertilizer types and two different locations, and the means \pm standard errors are provided.

The different fertilizer types had significantly influenced ($p < 0.001$) the plant height and emergence percentage compared to the control group but no difference among the treatment groups ($p > 0.05$). For instance, using sole vermi compost, a maximum height of 30.6 cm was recorded while the control group had the shortest plants, with a mean height of 21 cm. The results suggest that all fertilizer treatments positively influenced the plant height and emergence percentage of the carrot crop compared to the control group. The organic fertilizers showed comparable effects to the inorganic fertilizer on plant height and seed germination, indicating their potential as sustainable alternatives in carrot production. These findings align with previous research that demonstrated the positive impact of organic fertilizers on plant growth and health (Hati and Bandyopadhyay, 2011; Tadila, and Amare, 2019). Additionally, studies on the use of organic fertilizers in agriculture have highlighted their ability to improve soil fertility, nutrient availability, and overall crop productivity (Alzamel et al., 2022; Mahmud et al., 2016). The result of this study for carrot height (30.6 cm for vermicompost and 29.5 cm for conventional compost) was higher compared to the finding of Kifle and Birhanu (2019) who reported 29.16 cm by applying compost around Wolkite. However, the range of plant height in this study (16.5 cm to 35.4 cm) was lower than the findings of Amartey et al. (2022) who reported height ranging from 10 cm to 70 cm using different organic fertilizers. Similarly, the result of this study (30.6 cm for vermicompost) as lower in height compared to the finding of Afrin et al. (2019) who reported 43 cm by applying vermicompost. Overall, the use of organic fertilizers enhances the height and germination of carrot significantly.

Table 1. Emergence percentage and plant height of carrot under different fertilizers in Tehuledere district

Treatment	Plant Height (cm)			Emergence Percentage (%)		
	Gobeya	Libanos	Mean±SE	Gobeya	Libanos	Mean±SE
V compost	28.2 ^{bc}	33 ^c	30.6±0.8 ^b	92.6 ^c	94.1 ^c	93.4 ^b ±1.3
C compost	27.5 ^{bc}	31.6 ^c	29.5±0.8 ^b	94.2 ^c	93.3 ^c	93.8 ^b ±1.3
V and C mix	27.6 ^{bc}	32.8 ^c	30.2±0.8 ^b	92.9 ^c	95.1 ^c	94 ^b ±1.3
NPSB	31 ^{bc}	32.7 ^c	31.9±1.2 ^b	92.6 ^{bc}	91 ^c	91.8 ^b ±1.7
Control	23.1 ^{ab}	18.9 ^a	21± 0.8 ^a	83.1 ^{ab}	80.3 ^a	81.7 ^a ±1.3
Mean±SE	27.5 ^a ±0.6	29.8 ^b ±0.5	28.5	91.1 ^a ±1	90.8 ^a ±1	90.8
Range	16.5-35.4			76.8-98.6		
Interaction effect	**			NS		
SEM	1.15			1.6		
Treatment effect	***			***		

C=Conventional; V=Vermi; SE=standard error; SEM=standard error of means; ***=significantly different at p<0.001; **=significantly different at p<0.01; NS=not significant; Means with the same letter superscripts within same columns are not significantly different.

Leaf number and weight of carrot

Table 2 presents the results on leaf number, leaf fresh weight, and leaf dry weight in the two locations using the five treatment groups. The data is presented in terms of mean values and standard errors, as well as the range of values.

The fertilizers had varying effects on the aforementioned parameters. For instance, in leaf number, the highest was observed in the inorganic fertilizer (13 ± 0.8), while the control group (6.67 ± 0.5) had the lowest leaf number (p<0.001). The range values of leaf number (4.43 to 15.93) was far higher than the findings of Amartey et al. (2022) who reported a result ranging from 2 to 11 using different organic fertilizers. Similarly, the leaf number result of this study (12 for vermicompost) was higher compared to the finding of Afrin et al. (2019) who reported a leaf number of 9 via the application of vermicompost. However, the result of this study (12 for vermicompost and 10 for conventional compost) was lower in carrot leaf number compared to the finding of Kifle and Birhanu (2019) that reported 15 via application of 75 t/ha compost in Wolkite area. The inorganic fertilizer, NPSB, showed promising and statically similar results with the organic fertilizers like the vermi-compost in the study area; however, NPSB has negative environmental impact.

Table 2. Leaf number, leaf fresh weight and leaf dry weight of carrot under different fertilizers and two locations in Tehuledere district

Treatment	Leaf Number			Fresh leave weight (g)			Dry leave weight (g)		
	G	L	Mean±S E	G	L	Mean±S E	G	L	Mean±SE
V compost	12 ^c	12 ^c	12 ^c ±0.5	31.8 ^b	64 ^e	47.9 ^c ±1.8	8.4 ^{bcd}	6.77 ^{bcd}	7.58 ^b ±0.5
C compost	10.7 ^{bc}	9.33 ^{abc}	10 ^b ±0.5	20.5 ^{ab}	45.7 ^{cd}	33.1 ^b ±1.8	6.4 ^{bc}	7.8 ^{bcd}	7.12 ^b ±0.5
V and C mix	12 ^c	12 ^c	12 ^c ±0.5	25.6 ^b	49.9 ^d	37.8 ^b ±1.8	7.23 ^{bc}	9.43 ^{cd}	8.33 ^b ±0.5
NPSB	13 ^{bc}	13 ^c	13 ^c ±0.79	26 ^{abc}	58.3 ^{cdc}	42.2 ^{bc} ±3	7.62 ^{bc}	10.4 ^d	9.0 ^b ±0.8
Control	7.3 ^{ab}	6 ^a	6.67 ^a ±0.5	11.9 ^a	27.7 ^b	19.8 ^a ±1.8	2.37 ^a	5.53 ^{ab}	3.95 ^a ±0.5
Mean±SE	11 ^a ±0.4	11 ^a ±0.3	10.6	23 ^a ±2	49 ^b ±1	36.9	6.4 ^a	7.99 ^b	7.2
Range	4.43- 15.93			6.28-69.6			0.74- 12.02		
Interaction effect	NS			NS			*		
SEM	0.75			1.88			0.77		
Treatment effect	***			***			***		

G=Gobeya; L=Libanos; C=Conventional; V=Vermi; SE=standard error; SEM=standard error of means; ***=significantly different at p<0.001; *=significantly different at p<0.05; NS=not significant; Means with the same letter superscripts within same columns are not significantly different.

Yield and Yield Related Parameters of Carrot under Different Fertilizers and Locations

The effects of treatments on the yield and yield related parameter (root length, root diameter, root weight, marketable yield, unmarketable yield, and total yield) of carrot is indicated in table 3 and 4. The data are presented as means ± standard error and range.

All of the fertilizer types significantly improved (p<0.001) the yield of carrot compared to the control group. Vermi-compost resulted in the highest mean values (p<0.001) for root length, root diameter, root fresh weight, marketable yield, and total yield followed by the mixture of fertilizers, whereas the control had the lowest values for all parameters (Tables 3 and 4). The potentials of organic fertilizers particularly vermi-compost demonstrated improving the root yield of carrot. On the other hand, the use of the inorganic fertilizer, NPSB, provided comparable result with the vermi-compost (p>0.05), while it may result in negative impacts on soil health and environmental sustainability (Alzamel et al., 2022). These findings were consistent with the reports of Alzamel et al. (2022) and Mahmud et al. (2016) that have shown the beneficial effects of organic fertilizers on crop productivity, and have important implications for sustainable agriculture and the development of environmentally friendly farming practices. The study provides valuable information for farmers and researchers interested in sustainable and organic methods for improving carrot productivity. The result of this experiment indicated when

vermicompost is solely applied on a farmland; it played a significant role on the yield and yield contributing of carrot.

Length, diameter and fresh weight of carrot root

The root length values across all treatments ranged from 11.3 cm to 19.4 cm (Table 3). It is evident that the highest mean root length ($p < 0.001$) is observed in the "inorganic fertilizer" with a value of 21.3 whereas the lowest mean root length is recorded in the "control group" with a mean value of 12 cm. The other treatments, namely "sole vermi-compost", "conventional compost" and "mixture of conventional and vermi composts", fall in between with mean values of 19.6 cm, 17.1 cm, and 18.3 cm, respectively. The results of this study (19.8 cm and 17.1 cm) for sole vermicompost and for conventional compost, respectively for root length were in line with the findings of Kifle and Birhanu (2019) that reported 19.66 cm by applying conventional compost at a rate of 75t/ha in Wolkite area. However, the root length of this study (19.8 cm) for sole vermicompost was higher compared to the finding of Afrin et al. (2019) who reported a carrot root length of about 17 cm using vermicompost. Root length is a critical parameter as it indicates the extent of root development and exploration of the soil. Longer roots generally imply better nutrient and water absorption capabilities, which can lead to overall plant health and growth. The significantly higher root length observed in the "inorganic fertilizer" treatment suggests that this fertilizer might have provided better nutrient availability, promoting root growth despite its impact on pollution of the environmental.

The root diameter values across all treatments ranged from 1.35 cm to 6.95 cm (Table 3). The highest mean root diameter ($p < 0.001$) is observed in the treatment groups of "sole vermi-compost" and "inorganic fertilizer" with a mean value of 5.6 cm, while the lowest mean root diameter is recorded in the "control group" with a mean value of 2.27 cm. The result of this study (5.6 cm and 4.5 cm) for sole vermicompost and conventional compost, respectively for the root diameter was much higher than the findings of Kifle and Birhanu (2019) that reported 1.7 cm in Wolkite area. Similarly, the root length of this study (5.6 cm) for vermicompost was higher than the finding of Afrin et al. (2019) who reported a root length of about 3.5 cm via vermicompost application. Root diameter plays a crucial role in providing mechanical support to the plant and affects nutrient uptake efficiency. Generally, thicker roots indicate a stronger and

healthier root system. The significantly higher observed root diameters suggest that the treatment fertilizers may have contributed to root thickening and potentially improved structural support for the plants.

The range of fresh root weight values across all treatments ranged from 15.7 g/plant to 101.5 g/plant (Table 3). The highest mean fresh root weight (79.9 g/plant) was observed in the "sole vermi-compost" and "inorganic fertilizer" treatments. The lowest mean fresh root weight was recorded in the "control" with a value of 28.8 g/plant. The result of this study (79.9 g/plant for vermicompost and 57.6 g/plant for conventional compost) of fresh root weight was much higher than the findings of Kifle and Birhanu (2019) that reported 23.48 g/plant. However, the fresh root weight of this study (79.9 g/plant for vermicompost) was lower compared to the finding of Afrin et al. (2019) who reported a fresh root weight of about 91 g/plant using vermicompost. Fresh root weight is a critical indicator of plant growth and biomass production. A higher fresh root weight signifies increased root mass and, in turn, more nutrient and water absorption. The significantly higher fresh root weights in the treatments suggest that the fertilizers might have provided essential nutrients, leading to better root development and higher biomass production.

Table 3. Root length, root diameter and fresh root weight of carrot under different types of fertilizer in Tehuledere district

Treatment	Root length (cm)			Root diameter (cm)			Fresh root weight (g)		
	G	L	Mean±SE	G	L	Mean±SE	G	L	Mean±SE
V compost	19.4 ^{de}	19.8 ^{de}	19.6 ^c ±0.3	6.23 ^d	4.97 ^{cd}	5.6 ^c ±0.2	89.9 ^f	69.9 ^{de}	79.9 ^d ±2.25
C compost	17.1 ^b	17.1 ^b	17.1 ^b ±0.3	4.87 ^{cd}	4.17 ^{bc}	4.52 ^b ±0.2	68 ^{de}	47.3 ^{bc}	57.6 ^b ±2.25
V and C mix	18.1 ^{bc}	18.5 ^{bcd}	18.3 ^b ±0.3	5.37 ^{cd}	4.57 ^{cd}	4.97 ^{bc} ±0.2	78.7 ^{ef}	61.7 ^{cd}	70.2 ^c ±0.3
NPSB	20.2 ^{de}	20.5 ^{de}	21.3 ^d ±0.4	5.5 ^{cd}	5.7 ^{cd}	5.6 ^c ±3.6	89.9 ^{ef}	70 ^{de}	79.9 ^d ±3.2
Control	12.2 ^a	11.9 ^a	12 ^a ±0.3	2.47 ^{ab}	2.07 ^a	2.3 ^a ±0.2	35.5 ^{ab}	22.2 ^a	28.8 ^a ±2.25
Mean±SE	17.8 ^a ±0.3	17.6 ^a ±0.2	17.4	4.89 ^b ±0.2	4.29 ^a ±0.2	4.5	72.4 ^b ±1.8	54.2±1.6	61.4
Range	11.3-19.4			1.35-6.95			15.7-101.5		
Interaction effect	NS			NS			NS		
SEM	0.41			0.34			3.08		
Treatment effect	***			***			***		

G=Gobeya; L=Libanos; C=Conventional; V=Vermi; SE=standard error; SEM=standard error of means; ***=significantly different at p<0.001; NS=not significant; Means with the same letter superscripts within same columns are not significantly different.

Marketable, unmarketable, and total yields of carrot production

The result for the effects of fertilizer types on marketable yield, unmarketable yield, and total yield is presented in table 4. Such data provide valuable information on the effects of different fertilizer types on the productivity and sustainability of carrot in the study area.

Marketable yield refers to the quantity of crops that meet the quality standards required for sale or consumption. On the contrary, unmarketable yield refers to the quantity of crops that do not meet the required quality standards for sale or consumption. In this study, the highest marketable yield of carrot ($p < 0.001$) was found in the sole vermi-compost application ($41.7 \text{ t/ha} \pm 0.76$). Statically similar result ($p > 0.05$) was obtained in the inorganic fertilizer ($42.5 \text{ t/ha} \pm 1.14$) treatment group. On the other hand, the lowest mean marketable yield was recorded in the control group ($19 \text{ t/ha} \pm 0.76$). The marketable yield of this study (41.7 t/ha) for the sole vermicompost was found higher than the finding of Afrin et al. (2019) who reported a marketable yield of about 38 t/ha by applying vermicompost. Lower unmarketable yield is desirable, as it indicates less wastage and better overall crop quality. In this study, the control ($9.1 \text{ t/ha} \pm 0.45$) treatment had the highest unmarketable yield ($p < 0.001$) compared to the treatment groups.

Total yield is the sum of marketable and unmarketable yields, representing the overall productivity of the carrot. The vermi-compost ($45.5 \text{ t/ha} \pm 0.9$) and the inorganic fertilizer ($45.6 \text{ t/ha} \pm 0.9$) treatments again resulted in the highest total yields ($p < 0.001$). The lowest mean total yield was recorded in the control group ($28.9 \text{ t/ha} \pm 0.9$). The other treatments (conventional compost, and combination of vermi-conventional composts) were in between with a mean values of $36.9 \text{ t/ha} \pm 0.9$ and $41.4 \text{ t/ha} \pm 0.9$, respectively. The marketable carrot yield of this study (41.7 t/ha) for the sole vermicompost treatment was in line with the finding of Afrin et al. (2019) who reported a marketable yield of about 42 t/ha by applying sole vermicompost fertilizer. However, this study resulted in 24.1 t/ha to 52.1 t/ha which was higher than the findings of Amartey et al. (2022) who reported a total carrot yield ranging from 22.2 - 47.6 t/ha using different organic fertilizers.

Overall, the results suggest that the fertilizers (sole vermi-compost treatment and inorganic fertilizer) were the most effective in terms of yield. These treatments demonstrated better crop productivity compared to the conventional compost and the combination of fertilizers. The control group had the lowest marketable yield, indicating the necessity for appropriate agricultural practices to improve the carrot yield. The use of organic fertilizers not only enhance yield but also contributes to environmentally friendly farming practices and soil health improvement (Tadila and Amare, 2019). By adopting such practices, farmers in the study area can move towards more sustainable and eco-friendly agricultural methods.

Table 4. Marketable, unmarketable and total yields of carrot produced under different type of fertilizers in Tehuledere district

Treatment	Marketable yield (t/ha)			Unmarketable yield (t/ha)			Total yield (t/ha)		
	G	L	Mean±SE	G	L	Mean±SE	G	L	Mean±SE
V compost	44.3 ^c	39 ^c	41.7 ^d ±0.8	5 ^{abc}	2.77 ^a	3.88 ^a ±0.5	49 ^c	41.8 ^b	45.5 ^d ±0.9
C compost	33.1 ^b	30 ^b	31.5 ^b ±0.8	6.6 ^{bc}	4.03 ^{ab}	5.32 ^a ±0.5	40 ^b	34 ^c	36.9 ^b ±0.9
V and C mix	39.7 ^c	33.2 ^b	36.4 ^c ±0.8	6.4 ^{bc}	3.47 ^{ab}	4.95 ^a ±0.5	46 ^b	36.6 ^c	41.4 ^c ±0.9
NPSB	42 ^c	43 ^c	42.5 ^d ±1.1	4.2 ^{abc}	2.07 ^a	3.13 ^a ±0.7	46 ^b	45.1 ^b	45.6 ^d ±0.9
Control	20.5 ^a	19 ^a	19 ^a ±0.76	10.3 ^d	7.87 ^{cd}	9.1 ^b ±0.5	31 ^a	26.9 ^a	28.9 ^a ±0.9
Mean±SE	35.9 ^b ±0.6	32.8 ^a ±0.5	33.8	6.5 ^b ± 0.4	4.04 ^a ±0.3	5.4	42 ^a ±0.6	36.9 ^a ±0.6	39.2
Range	16.7-46.6			0.751-11.65			24.1-52.1		
Interaction effect	NS			NS			NS		
SEM	1.07			0.66			1.33		
Treatment effect	***			***			***		

G=Gobeya; L=Libanos; C=Conventional; V=Vermi; SE=standard error; SEM=standard error of means; ***=significantly different at p<0.00; NS=not significant; Means with the same letter superscripts within same columns are not significantly different.

Conclusion

The result of the experiment indicated that applications of the fertilizers in general played an important role and the potential of the organic fertilizers in particular are demonstrated in improving the growth and yield of carrot. Applications of the sole vermi-compost and inorganic fertilizer (NPSB) were found the most effective having better and comparable results. However, the applications of inorganic fertilizers like NPSB have negative environmental impacts leading to unsustainable agricultural production. As a result, sole application of vermi-compost is recommended for farmers of Tehuledere district and similar agroecology for improving carrot productivity in organic methods while contributing for sustainable agriculture.

Acknowledgement

We acknowledged extension workers of Gobeya and Libanos *kebeles* for their field support and data collection.

Funding

This research project is funded by Swiss Agency for Development and Cooperation (SDC) in collaboration with Biovision Africa Trust.

Data Availability Statement

Data are available from the corresponding author on reasonable request.

Consent for publication

The authors have provided their consent for publication.

Competing interests

The authors declare that they have no competing interests.

Author Contribution

The first four authors (Faris Hailu, Seid Hassen, Seid Hussien, and Eshetu Belete) had equal contribution involved in the whole processes of the project starting from proposal development. The last author, Tewodros Alemu, had participated in data analysis and manuscript writes up.

References

- Afrin, A., Islam, M.A., Hossain, M.M. and Hafiz, M.M.H. 2019. Growth and yield of carrot influenced by organic and inorganic fertilizers with irrigation interval. *Journal of Bangladesh Agricultural University*, 17(3): 338–343. <https://doi.org/10.3329/jbau.v17i3.43207>
- Alzamel, Nurah M., Eman M. M. Taha, Abeer A. A. Bakr, and Naglaa Loutfy. 2022. "Effect of Organic and Inorganic Fertilizers on Soil Properties, Growth Yield, and Physiochemical Properties of Sunflower Seeds and Oils" *Sustainability* 14, no. 19: 12928. <https://doi.org/10.3390/su141912928>
- Amartey, J. N. A., Sarkordie-Addo, J., Essilfie, M. E., & Dapaah, H. K. (2022). Growth and yield of carrots affected by integrated nutrient management of organic and inorganic fertilizers. *African Journal of Agricultural Research*, 18(7), 576-585.
- Basel Natsheh and Sami Mousa, 2014. Effect of Organic and Inorganic Fertilizers Application on Soil and Cucumber (*Cucumis Sativa L.*) Plant Productivity, *International Journal of Agriculture and Forestry*, Vol. 4 No. 3, 2014, pp. 166-170. doi: 10.5923/j.ijaf.20140403.03.
- Ellis, R.H. and Roberts, E.H. (1980) Improved Equations for the Prediction of Seed Longevity. *Annals of Botany*, 45, 13-30. <https://doi.org/10.1093/oxfordjournals.aob.a085797>
- Getachew, T. and Mohammed, Y. (2012). Mapping the Current Knowledge of Carrot Cultivation in Ethiopia. Technical Report Submitted to Carrot Aid, Charlottenlund, Denmark.
- Hati, K. and Bandyopadhyay, K. (2011). Fertilizers (Mineral, Organic), Effect On Soil Physical Properties. In: Gliński, J., Horabik, J., Lipiec, J. (eds) *Encyclopedia of Agrophysics*. *Encyclopedia of Earth Sciences Series*. Springer, Dordrecht. https://doi.org/10.1007/978-90-481-3585-1_201
- Jamuna K.S., Ramesh C.K., Srinivasa T.R., Raghu K.L., 2011. In vitro antioxidant studies in some common fruits. *International Journal of Pharmacy and Pharmaceutical Sciences* 3.1 (2011) 60-63.

Kifle Zerga and Birhanu Tsegaye, 2019. Effect of Different Rates of Compost Application on Growth Performance and Yield Components of Carrot (*Daucus carota* L.) in Gurage Zone, Ethiopia. *International Journal of African and Asian Studies*, 54, 24-31.

Mahmud AJ, Shamsuddoha ATM, Haque MN, 2016. Effect of Organic and Inorganic Fertilizer on the Growth and Yield of Rice (*Oryza sativa* L). *Nat Sci* 2016;14(2):45-54. doi:10.7537/marsnsj14021607.

Nambia V.S., Daniel M., Guin P. 2010. Characterization of Polyphenols from Coriander leaves (*coriandrum sativum*), red amaranthus (*a. paniculatus*) and green amaranthus (*a. frumentaceus*) using paper chromatography: and their health implications. *Journal of Herbal Medicine and Toxicology*, 4: 173-177

Shashi Kamal, Mohit Kumar*, Rajkumar and Manoj Raghav, 2018. Effect of Biofertilizers on Growth and Yield of Tomato (*Lycopersicon esculentum* Mill). *Int.J.Curr.Microbiol.App.Sci* (2018) 7(2): 2542-2545

Tadila Getaneh, and Amare Mezgebu. (2019). The effect of different rates of Vermicompost and Inorganic Fertilizers on Growth and Yield of Cabbage (*Brassica oleracea* L.) at Mojo in Ethiopia. *Journal of Soil Science and Environmental Management*, 10(8), 99-105.